

## **WEEKLY NEW BRUNSWICK INFLUENZA REPORT**

Reporting period: January 20 to January 26 2013 (week 4)

### **Summary**

#### **In New Brunswick, positive detections remain high and increases in ILI consultation and number of reported outbreaks**

##### **New Brunswick:**

- There have been 319 positive influenza detections during week 4, 318 influenza A (unsubtyped) and 1 influenza B.
- The ILI consultation rate increased and was above the expected range for this time of year.
- An increase of ILI and influenza-confirmed outbreaks were reported, 3 in Nursing Homes and 37 in schools.

##### **Canada:**

- Several indicators of influenza activity decreased; however, many regions across Canada continue to report widespread and localized influenza activity.
- 1919 laboratory detections of influenza were reported, proportion of positive tests was 22.3%.
- The ILI consultation rate was above the expected level for this time of year. 104 influenza outbreaks were reported, 31 in long-term care facilities, 6 in hospitals, 45 in schools and 22 in other settings.

##### **International:**


- United States: From July 12 through January 26, 2013, a total of 312 infections with variant influenza viruses (308 H3N2v viruses, 3 H1N2v viruses and 1 H1N1v virus) have been reported in 11 states.

### **1) Influenza Laboratory Data<sup>1</sup>**

- Percent positive laboratory results remain high in week 4.
- 319 influenza detections were reported during that period.
- Since the beginning of the season, 765 positive influenza detections have been reported, 7 influenza A (H1N1)pdm09, 379 influenza A (H3) viruses, 376 influenza A (unsubtyped) and 3 influenza B.

<sup>1</sup> Surveillance specimens are submitted by recruited New Brunswick Sentinel Practitioner Influenza Network (NB SPIN) practitioners, which are comprised of 8 sites in Emergency Rooms, 5 sites in Family Practice, 3 sites in First Nations communities, 1 site in a Nursing Home, 3 sites in Universities and 8 sites in Community Health Centers. Diagnostic specimens are submitted by physicians in the community/hospital setting. Influenza laboratory data is comprised of results from surveillance and diagnostic specimens. All laboratory specimens are tested using a real-time PCR assay, which is a rapid detection method designed for detection of all known variants of influenza A and B. All laboratory-confirmed cases are reported for the week when laboratory confirmation was received.

**Graph 1:** Number and percent of positive influenza specimens in New Brunswick, by week, up to January 26, 2013 (data source: G. Dumont lab results)



**Table 1: Positive influenza test results by Health Region in New Brunswick up to January 26, 2013 (data source: G. Dumont lab results)**


Region	Reporting period:						Cumulative: (2012/2013 season)					Cumulative: (2011/2012 season)		
	January/20/2012–January/26/2013						Aug./26/2012 –Jan./26/2013					Aug./28/2011 – Aug./25/2012		
	Activity level <sup>2</sup>	A				B	A				B	A		B
A(H1)		A(H3)	(H1N1) pdm09	unsubt yped		A(H1)	A(H3)	(H1N1) pdm09	unsubt yped		Non-(H1N1) pdm09	(H1N1) pdm09		
Region 1	Localized	0	0	0	127	0	0	129	6	152	2	24	17	206
Region 2	Localized	0	0	0	43	1	0	30	1	52	1	1	2	18
Region 3	Localized	0	0	0	17	0	0	21	0	21	0	1	0	3
Region 4	Sporadic	0	0	0	44	0	0	33	0	52	0	2	9	2
Region 5	Localized	0	0	0	3	0	0	10	0	3	0	0	2	4
Region 6	Localized	0	0	0	67	0	0	132	0	76	0	1	6	16
Region 7	Localized	0	0	0	17	0	0	24	0	20	0	1	1	12
Total NB		0	0	0	318	1	0	379	7	376	3	30	37	259

<sup>2</sup> Influenza activity level definition is available on the PHAC FluWatch website: <http://www.phac-aspc.gc.ca/fluwatch/12-13/def12-13-eng.php>

## 2) ILI Consultation Rates<sup>3</sup>

- During week 4, the ILI consultation rate was 65.5 consultations per 1,000 patient visits, and was above the expected levels for this time of year.
- During week 4, the sentinel response rate was 32% for both the FluWatch sentinel physicians and the NB SPIN practitioners.

Graph 2: ILI Consultation Rates in New Brunswick, by report week, season 2012/13 compared to previous seasons\*



\* The mean rate was based on data from the 1996/97 to 2011/2012 seasons and excludes the Pandemic season (2009-2010).

<sup>3</sup> A total of 38 practitioner sites (20 FluWatch sentinel physicians and 18 NB SPIN sites) are recruited this season to report the number of ILI patients and total patient consultations one day during a reporting week.

### 3) ILI and Laboratory-Confirmed Outbreak Data


Table 2: ILI activity/outbreaks in New Brunswick nursing homes and schools for the reporting week, current and previous seasons.

	Reporting period: January/20/2012 –January/26/2013			Cumulative # of outbreaks season 2012-2013	Cumulative # of outbreaks season 2011-2012
	Lab-confirmed outbreaks in Nursing Homes*	Schools reporting ILI outbreaks**	Lab-confirmed outbreaks in Other Settings*		
Region 1	0 out of 13	3 out of 74	0	4	4
Region 2	2 out of 15	16 out of 81	0	22	6
Region 3	0 out of 14	6 out of 95	0	11	8
Region 4	0 out of 6	0 out of 22	0	1	2
Region 5	0 out of 2	1 out of 18	0	3	7
Region 6	0 out of 9	8 out of 35	0	15	2
Region 7	1 out of 4	3 out of 27	0	6	2
Total NB	3 out of 63	37 out of 352	0	62	31

\*Two or more ILI cases within a seven day period, including at least one laboratory-confirmed case of influenza. Outbreaks are reported in the week when laboratory confirmation is received.

\*\*Schools reporting greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or Public Health Authority) which is likely due to ILI.

**Graph 3: Number of Influenza Outbreaks in Nursing Homes<sup>1</sup> and ILI Outbreaks in Schools<sup>2</sup> reported to Public Health in New Brunswick, by report week, season 2012/13.**



<sup>1</sup> The National FluWatch definition of an outbreak in a nursing home is stated as two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case.

<sup>2</sup> The National FluWatch definition of an ILI outbreak in a school is stated as absenteeism greater than 10% (or absenteeism that is higher (e.g.>5-10%) than expected level as determined by school or Public Health Authority) which is likely due to ILI.

**National Flu Watch Program** - Additional information on influenza activity in Canada and around the world is available on the Public Health Agency of Canada’s website at:

[www.phac-aspc.gc.ca/fluwatch/index.html](http://www.phac-aspc.gc.ca/fluwatch/index.html)

**Other Links:**

World-

[http://www.who.int/influenza/surveillance\\_monitoring/updates/latest\\_update\\_GIP\\_surveillance/en/index.html](http://www.who.int/influenza/surveillance_monitoring/updates/latest_update_GIP_surveillance/en/index.html)

Europe: [http://www.euroflu.org/cgi-files/bulletin\\_v2.cgi](http://www.euroflu.org/cgi-files/bulletin_v2.cgi) and

[http://www.ecdc.europa.eu/en/healthtopics/seasonal\\_influenza/epidemiological\\_data/Pages/Weekly\\_Influenza\\_Surveillance\\_Overview.aspx](http://www.ecdc.europa.eu/en/healthtopics/seasonal_influenza/epidemiological_data/Pages/Weekly_Influenza_Surveillance_Overview.aspx)

PAHO: [http://new.paho.org/hq/index.php?option=com\\_content&task=blogcategory&id=805&Itemid=569](http://new.paho.org/hq/index.php?option=com_content&task=blogcategory&id=805&Itemid=569)

Australia: <http://www.health.gov.au/internet/main/publishing.nsf/Content/cda-surveil-ozflu-flucurr.htm>

New Zealand: [http://www.surv.esr.cri.nz/virology/influenza\\_weekly\\_update.php](http://www.surv.esr.cri.nz/virology/influenza_weekly_update.php)

Argentina: : <http://www.msal.gov.ar/>

South Africa: <http://www.nicd.ac.za/>

US: [www.cdc.gov/flu/weekly/](http://www.cdc.gov/flu/weekly/)